

Vector Solitons and Spontaneous Lorentz Violation Mechanism

L.P. Colatto*

LAFEX, CBPF, Rua Xavier Sigaud, 150, CEP 22290-180, Rio de Janeiro, RJ, Brasil

A.L.A. Penna[†] and W.C. Santos[‡]

IF, Universidade de Brasília, CEP 70919-970, Brasília, DF, Brasil

(Dated: February 2, 2008)

Taking a full 3D nonlinear vector matter field dynamics, a vector version of a soliton state was found. The Nielsen-Olesen procedure was used in order to derive a Lorentz-violation vector parameter which characterizes, via Spontaneous Symmetry Breaking mechanism, the non-trivial vacuum. A stable vortex configuration is obtained, and although the Chern-Simons-type terms do not contribute to the value of the vortex core, the propagator analysis suggests us the possibility of a contribution to the size of the vortex core and to the growth of the field to achieve the asymptotic limit value with the distance.

PACS numbers:

I. INTRODUCTION

Since the birth of quantum mechanics there has been a continuous growing interest in the study of non-commutative possible phenomena [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12], and its consequent (or not) Lorentz violation effects. In the context of field theory it can be determined by introducing deformation terms in the algebra of the objects of the model or in its own dynamics, particularly when we are dealing with interacting models. In the 90's the q-deformation introduced to algebras and the quantum groups developed an interesting theoretical scenario [13]. From another point of view, Colladay and Kostelecký [14, 15], have observed the possible occurrence of Lorentz violating extensions to the Standard Model. Since their pioneering work, the Lorentz symmetry violation has been extensively investigated with the strings theory as the underlying motivation. Indeed, some astrophysical phenomena could be fitted by these extensions [16], and theoretical developments have been largely worked on [17, 18, 19].

An interesting way to theoretically represent these phenomena is through the vacuum of a model with Chern-Simons photon coupling, what we have called Carroll-Jackiw field model [20]. This work has presented a preferred space-time direction dictated by the introduction of a constant four-vector b_μ . Consequently, the photon circular polarization shows a non usual dispersion relation [21], while the linear polarized photons exhibit optical birefringence. Furthermore the four-vector induces anisotropic optical vacuum which could be detected through the observation of tiny Lorentz-breaking effects ($b_\mu b^\mu < 10^{-33} \text{eV} \simeq 10^{-28} \text{cm}^{-1}$) on space-time [22, 23, 24, 25]. Indeed, a nontrivial vacuum value indicates that a fundamental symmetry-breaking occurs for such “field” at low-energy limit. A careful treatment of this issue suggests that “spontaneous” Lorentz-violation terms, with consequent anisotropic optical effects, could originate from a spontaneous symmetry breaking (SSB) mechanism on a matter vector field [26]. Taking this into account, the reduced model to $(1+2)D$ may reveal the contribution of topological objects to such nontrivial optical vacuum effects. Our purpose in this work is to obtain stable $(1+2)D$ vortex-lines solutions starting from interacting nonlinear matter vector field Lagrangian, and also to characterize the vector nature of these topological objects. We are going to discuss the contribution of the Chern-Simons-type terms to asymptotic behavior of the matter field.

The outline of the work is the following: In Sec. II we consider a nonlinear matter vector field model and we discuss how it can arise in stable formation for vector solitons. Sec. III is devoted to computing the vortex-lines via Nielsen-Olesen way and verifying the possible solutions and stability. In Sec. IV we make a study of the mass poles through the propagators obtained and discuss the relations to the vacuum and asymptotic limit of the fields. Finally, in the Conclusion we discuss the results and give perspectives of new approaches to the subject.

*Electronic address: colatto@cbpf.br

†Electronic address: andrel@fis.unb.br

‡Electronic address: wytler@fis.unb.br

II. VORTEX LINES IN THE MATTER VECTOR FIELD MODEL

Taking the reference [26], we recall the charged matter vector field model and we assume it as living in a $(1+2)D$ world, including global $U(1)$ invariant terms, and topological terms. It can be written as,

$$\mathcal{L} = -\frac{1}{2}F_{\mu\nu}^*F^{\mu\nu} - (\partial_\mu B^\mu)^*(\partial_\nu B^\nu) + m\epsilon^{\mu\nu\kappa}B_\mu\partial_\nu B_\kappa^* + m\epsilon^{\mu\nu\kappa}B_\mu^*\partial_\nu B_\kappa + \alpha^2 B_\mu^*B^\mu + \lambda(B_\mu^*B^\mu)^2 \quad (1)$$

where B_μ is a charged vector field and $F_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$ is the field strength. The symmetry $U(1)$ allows us to insert a topological massive Chern-Simons-type term that does not play a role in the vacuum achievement but will be important to the mass term definition in the asymptotic limit behavior that will be made clear further ahead¹. As we are treating B_μ as a matter field the global $U(1)$ symmetry induces the presence of a dynamical longitudinal term in the Lagrangian (1). We can verify that this model gives rise to a SSB mechanism whose non-trivial vacuum [17, 18, 26] allows verification of particular soliton solution. Analogously to the usual scalar case we are going to assume asymptotic values on the boundary space for an S^1 topological circle recalling the Nielsen-Olesen approach. Consequently, taking r and θ as the polar coordinates, we can construct the following values for B_μ at $r \rightarrow \infty$,

$$B_\mu = b_\mu e^{in\theta} \quad \text{and} \quad B_\mu^*B^\mu = b_\mu b^\mu = b^2 \quad (2)$$

where b_μ is a constant vector that is defined by a minimal value of the energy functional in Lagrangian (1), as it also has been assumed in [26]. The result is that the Lagrangian (1) allows a SSB on matter vector field B_μ to $\alpha^2 < 0$ and we have that,

$$b_\mu = \sqrt{\frac{-\alpha^2}{2\lambda}}u_\mu, \quad (3)$$

where u_μ is an unitary vector defined on the Minkowski space in $(1+2)D$ along the b_μ direction. Hence our particular choices for B_μ at infinity lead us to a fixed and exact value for the degenerated vacuum. Another interesting aspect in SSB mechanism applied to the matter vector field are the Lorentz-violation parameters generated spontaneously too. Indeed, as expected, the Lorentz-breaking term, b_μ -vector, occurs due to the original vector-nature of the lowest energy field [26].

We can observe that vortex lines structures can be derived from the non-trivial topology of the vacuum of the matter vector field. From the anisotropic vacuum in $(1+2)D$, the time-component ($\mu = 0$) and the space-component ($\mu = i$) are written as,

$$B_0 = b_0 e^{in\theta}, \quad B_i = b_i e^{in\theta}, \quad (4)$$

where the r and θ components in the plane are signaled by

$$B_r = b_r e^{in\theta} \quad B_\theta = b_\theta e^{in\theta} \quad (5)$$

where the b_0 , b_i (b_r and b_θ) are time, vector (radial and angular) components of the Lorentz-violating parameter. Several recent discussions related to the existence of these parameters have been offered [17, 18, 24, 25]. In this case, the B_μ field generates a vector soliton with arbitrary anisotropic dependence whose stability is analyzed starting from static configuration for the Hamiltonian functional density,

$$\mathcal{H}_{s.c.} = -(\vec{\nabla}B_0)^*(\vec{\nabla}B_0) + (\vec{\nabla} \times \vec{B})^*(\vec{\nabla} \times \vec{B}) + (\vec{\nabla} \cdot \vec{B})^*(\vec{\nabla} \cdot \vec{B}), \quad (6)$$

where we have also admitted a static configuration for the SSB potential,

$$V(B_\mu, B_\mu^*) = \lambda[b^2 - B_\mu^*B^\mu]^2 = 0, \quad (7)$$

which results precisely in the expression (2) on the boundary. We have also verified that the topological term in (1) does not contribute to the Hamiltonian density. Let us now treat the system (6) in three dimensions, admitting a cylindric or axial symmetry around the z -axis. Thus, the z -component of B_μ is assumed to be constant, and from (2) we have found that,

$$\vec{\nabla}B_0 = \frac{inb_0}{r}e^{in\theta}\hat{\theta}, \quad \vec{\nabla} \cdot \vec{B} = \frac{1}{r}(b_r + inb_\theta)e^{in\theta}, \quad \vec{\nabla} \times \vec{B} = \frac{1}{r}(b_\theta - inb_r)e^{in\theta}\hat{z}. \quad (8)$$

¹ The two Chern-Simons-type term are indeed symmetric, we have written them to observe the global $U(1)$ -symmetry explicitly.

Hence substituting (8) with (6) we find that at $r \rightarrow \infty$ the Hamiltonian functional density is given by,

$$\mathcal{H}_{s.c.} = \frac{n^2 b_0^2 + (n^2 + 1)(b_r^2 + b_\theta^2)}{r^2} \quad (9)$$

and the energy of the vector soliton relative to the anisotropic vacuum parameters is,

$$E = \int_0^\infty \mathcal{H}_{s.c.} r dr d\theta = [n^2 b_0^2 + (n^2 + 1)(b_r^2 + b_\theta^2)] \ln |r| \Big|_0^\infty \quad (10)$$

On the other hand, as in the case of the scalar theories, the vector soliton model also presents a logarithmic divergence. In a similar way, this can be treated by adding a gauge field A_μ in the system,

$$D_\mu B_\nu = \partial_\mu B_\nu + ie A_\mu B_\nu. \quad (11)$$

We shall then assume the gauge choice $A_0 = 0$, where $A_\mu = A = \frac{1}{e} \vec{\nabla}(n\theta)$, for very large r value. As is well known, we can write it analytically in cylindrical symmetry, or

$$A_r \rightarrow 0 \quad \text{and} \quad A_\theta \rightarrow -\frac{n}{er} \quad (12)$$

since we treat the covariant derivative (11) at infinity, the results is

$$D_\mu B_\nu = 0. \quad (13)$$

Thus, the stability of this model requires a “vector electrodynamics” whose Lagrangian model is defined by,

$$\mathcal{L} = -\frac{1}{4} f_{\mu\nu} f^{\mu\nu} - \frac{1}{2} \mathcal{F}_{\mu\nu}^* \mathcal{F}^{\mu\nu} - (D_\mu B^\mu)^* (D_\nu B^\nu) + m\epsilon^{\mu\nu\kappa} B_\mu^* (D_\nu B_\kappa) + m\epsilon^{\mu\nu\kappa} B_\mu (D_\nu B_\kappa)^* + \lambda[b^2 - B_\mu^* B^\mu]^2 \quad (14)$$

where $\mathcal{F}_{\mu\nu} = D_\mu B_\nu - D_\nu B_\mu$ and $f_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. From the relations (12) which define a pure gauge, the Lagrangian model (14) assumes a finite energy configuration for its Hamiltonian functional density at infinity,

$$\mathcal{F}_{\mu\nu} \rightarrow 0 \quad \text{and} \quad \mathcal{H} \rightarrow 0 \quad (15)$$

and the gauge field A_μ gives rise to a soliton magnetic flux,

$$\Phi = \oint \vec{A} \cdot d\vec{l} = -\frac{2\pi n}{e}, \quad (16)$$

which is equivalent to the result obtained for scalar field model and could be seen as magnetic flux quantization as in the London and Ginzburg-Landau equations to describe the type-II superconductors [27].

III. VECTOR MODEL FOR THE NIELSEN-OLESEN EQUATIONS

Based on the Lagrangian (14) we now consider a vector version to the Nielsen-Olesen equations. A remarkable feature of this study is the existence of a non-trivial vacuum for the B_μ field, which naturally guarantees the appearance of an anisotropic parameter via SSB mechanism. In this way a vector soliton object with non-zero spin emerges from this model whose dynamical equations depend on the quadratic norm of the anisotropic vector b_μ . We then assume the Lagrangian model

$$\mathcal{L} = -\frac{1}{4} f_{\mu\nu} f^{\mu\nu} - \frac{1}{2} \mathcal{F}_{\mu\nu}^* \mathcal{F}^{\mu\nu} - (D_\mu B^\mu)^* (D_\nu B^\nu) + m\epsilon^{\mu\nu\kappa} B_\mu^* (D_\nu B_\kappa) + m\epsilon^{\mu\nu\kappa} B_\mu (D_\nu B_\kappa)^* + \alpha^2 B_\mu^* B^\mu + \lambda(B_\mu^* B^\mu)^2 \quad (17)$$

whose equations of motion for the model are written as,

$$ie(B_\nu^* F^{\mu\nu} - B_\nu F^{\mu\nu*}) + ie(B^{\mu*} \partial_\nu B^\nu - B^\mu \partial_\nu B^{\nu*}) - iem\epsilon^{\mu\nu\kappa} (B_\nu^* B_\kappa - B_\kappa B_\nu^*) - 2e^2 A^\mu (B_\nu^* B^\nu) = \partial_\nu f^{\mu\nu}, \quad (18)$$

$$D_\mu D^\mu B^\nu + 2m\epsilon^{\kappa\lambda\nu} D_\lambda B_\kappa = [\alpha^2 + 2\lambda(B_\mu^* B^\mu)] B^\nu. \quad (19)$$

Now we seek vortex solutions in the system described by (18). In order to extract further information about the system we admit that

$$A_\theta(r) = A(r) \quad \text{and} \quad B_\mu = \beta_\mu(r)e^{in\theta}, \quad (20)$$

where the limits to $\beta_\mu(r)$ are given by,

$$\lim_{r \rightarrow \infty} \beta_\mu(r) = b_\mu \quad \lim_{r \rightarrow 0} \beta_\mu(r) = 0, \quad (21)$$

here, $B_\mu^* B^\mu = |\beta(r)|^2$ and $|\beta(r)|^2 \rightarrow (|\beta(r)|_\infty)^2 = b^2$ at infinity. In the Minkowski space $(1+2)D$ we can observe that,

$$|\beta(r)| = \sqrt{\beta_0(r)^2 - \beta_r(r)^2 - \beta_\theta(r)^2} \quad \text{and so} \quad |\beta(r)|_\infty = |b| = \sqrt{b_0^2 - b_r^2 - b_\theta^2}. \quad (22)$$

Then the equations of motion (18) and (19) assume, in the static configuration, the forms:

$$\begin{aligned} \partial_i f^{\mu i} = & ie[B_0^* \partial^\mu B^0 + B_i^* \partial^\mu B^i - B_i^* \partial^i B^\mu - B_0 \partial^\mu B^{0*} - B_i \partial^\mu B^{i*} + B_i \partial^i B^{\mu*}] \\ & + ie[B^{\mu*} \partial_i B^i - B^\mu \partial_i B^{i*}] + ime\epsilon^{\mu\nu\kappa}(B_\nu^* B_\kappa - B_\nu B_\kappa^*) - 2e^2 A^\mu B_0 B^0 - 2e^2 A^\mu B_i^* B^i, \end{aligned} \quad (23)$$

$$D_i D^i B^\nu + 2m\epsilon^{\kappa i \nu} D_i B_\kappa = [\alpha^2 + 2\lambda(B_\mu^* B^\mu)] B^\nu. \quad (24)$$

Then by using the relations (20) and (21) in cylindric coordinates the equations of motion, (23) and (24), can be written down as,

$$\frac{d^2 A(r)}{dr^2} + \frac{1}{r^2} \frac{dA(r)}{dr} - \left[e^2 |\beta(r)|^2 + \frac{1}{r} \right] A(r) - \frac{2en}{r} |\beta(r)|^2 = 0, \quad (25)$$

$$\frac{1}{r} \frac{d}{dr} \left[r \frac{d\beta_0(r)}{dr} \right] - \left[\left(\frac{n}{r} - eA(r) \right)^2 + \alpha^2 + 2\lambda |\beta(r)|^2 \right] \beta_0(r) = 2m \left[-\frac{d\beta_\theta(r)}{dr} + \left(\frac{in}{r} + ieA(r) \right) \beta_r(r) \right], \quad (26)$$

$$\frac{1}{r} \frac{d}{dr} \left[r \frac{d\beta_r(r)}{dr} \right] - \left[\left(\frac{n}{r} - eA(r) \right)^2 + \alpha^2 + 2\lambda |\beta(r)|^2 \right] \beta_r(r) = 2m \left[\frac{in}{r} + ieA(r) \right] \beta_0(r), \quad (27)$$

$$\frac{1}{r} \frac{d}{dr} \left[r \frac{d\beta_\theta(r)}{dr} \right] - \left[\left(\frac{n}{r} - eA(r) \right)^2 + \alpha^2 + 2\lambda |\beta(r)|^2 \right] \beta_\theta(r) = -2m \frac{d\beta_0(r)}{dr}. \quad (28)$$

Unfortunately the analytical solution of these four equations can not be obtained. Recalling the scalar case, we have two, simpler equations but without analytical solution all the same [27]. In the above equations one simple particular solution can be obtained by making $|\beta(r)|$ for large r achieve the particular expressed in Eq. (22).

We can observe that the equation (25) is the consequence of static configuration when we substitute the $\beta_\mu(r)$ function into the equation of motion (23). The expressions (26), (27) and (28) are the components of the equation of motion (24) to $\mu = 0, r, \theta$ respectively. The solution of these coupled equations of motion system reveal the characteristic functions to each component $\beta_0(r), \beta_r(r), \beta_\theta(r)$. Unfortunately, again, this system can not admit an exact and general analytic solution. The only particular analytic solution we can obtain is through values of $|\beta(r)|^2$ near infinity, which implies that

$$|\beta(r)|^2 \simeq b^2. \quad (29)$$

In this case, we have that $|b|^2 = b^2$, so we can define the gauge field $A(r)$ and the magnetic field $H(r)$ are the θ - and z -component of their respective fields, or

$$\begin{aligned} A(r) \equiv A_\theta(r) &= A, \\ H(r) \equiv H_z(r) &= \frac{1}{r} \frac{d[rA(r)]}{dr}. \end{aligned} \quad (30)$$

Allowing these conditions, we obtain the following solutions to the fields as functions on the breaking parameter $|b|$,

$$\begin{aligned} A(r) &= -\frac{n}{er} - \frac{|b|}{e} K_1(e|b|r) \Big|_{r \rightarrow \infty} = -\frac{n}{er} + |b| \sqrt{\frac{\pi}{2er}} \exp(-e|b|r) + \mathcal{O}\left(\frac{1}{r}\right), \\ H(r) &= \frac{|b|}{e} K_0(e|b|r) \Big|_{r \rightarrow \infty} = |b| \sqrt{\frac{\pi|b|}{2er}} \exp(-e|b|r) + \mathcal{O}\left(\frac{1}{r}\right), \end{aligned} \quad (31)$$

where $K_0(e|b|r)$ and $K_1(e|b|r)$ are the modified Bessel functions, whose values are taken at infinity. We notice that in the above solutions, $|b|$ is a parameter of $A(r)$ and $H(r)$ that imposes a specific direction in space-time and consequently breaks the Lorentz symmetry². So, we can state that the magnetic field $H(r)$ is generated by a Lorentz symmetry violation. We also observe that, in spite of the absence of an exact solution to the coupled system of equations we can infer the solutions of the parameters $\beta_0(r)$, $\beta_r(r)$, $\beta_\theta(r)$ in good agreement with functional behavior when the system is going to infinity. In this limit we can suggest that the variation of these parameters is generically described through the equation,

$$\beta_\mu(r) = b_\mu \left(1 - ae^{-\sqrt{-\rho^2}r}\right), \quad (32)$$

where a and $-\rho^2$ are two real factors to be determined. The a -factor can be easily obtained ensuring that the solution obeys the asymptotic behavior. On the other hand, the ρ -factor is related to the mass of the field. Bearing this in mind and with the presence of a topological term in the original Lagrangian both might influence the vortex structure. To this end we use the propagator computation derived from the Lagrangian (1). We verify that close to the vortex core there is no contribution from the topological Chern-Simons-type term, but it is relevant to the process to reach the asymptotic limit, which can be dictated by a mass term parameter that will be obtained by a spectrum analysis.

IV. THE SPECTRUM ANALYSIS AND THE ASYMPTOTIC LIMIT BEHAVIOR

Taking the Lagrangian (1), the propagators can be written down in three sectors, as

$$\Omega^{\mu\nu} = \frac{\partial^\mu \partial^\nu}{\square}, \quad \Theta^{\mu\nu} = \eta^{\mu\nu} - \frac{\partial^\mu \partial^\nu}{\square}, \quad S^{\mu\nu} = \epsilon^{\mu\lambda\nu} \partial_\lambda, \quad (33)$$

where Ω is the longitudinal, Θ and S are the transversal sectors of the propagator remarking that S has topological origin. Thereby the invertible operator is $\mathcal{O}^{\mu\nu} = 2mS^{\mu\nu} + (\square + \alpha^2)\eta^{\mu\nu}$, and its inverse $\mathcal{O}^{-1}$ is

$$(\mathcal{O}^{\mu\nu})^{-1} = -\frac{1}{(\square + \alpha^2)} \Omega_{\mu\nu} + \frac{\square + \alpha^2}{(\square + \alpha^2)^2 + 4m^2 \square} \Theta_{\mu\nu} - \frac{2m}{(\square + \alpha^2)^2 + 4m^2 \square} S_{\mu\nu}. \quad (34)$$

Converting to the momenta space, the B -field propagator can be written down as ,

$$\langle B^\mu B^\nu \rangle_L(k) = -\frac{i}{k^2 - \alpha^2} \left(\frac{k^\mu k^\nu}{k^2} \right), \quad (35)$$

$$\langle B^\mu B^\nu \rangle_T(k) = \frac{i}{(k^2 - \alpha^2)^2 - 4m^2 k^2} \left[(k^2 - \alpha^2) \left(\eta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right) + 2mi\epsilon^{\mu\lambda\nu} k_\lambda \right], \quad (36)$$

where we have separated in longitudinal (L) and transversal (T) sectors. Observe that the longitudinal sector exhibits a simple pole $k^2 = \alpha^2$, and if we saturate this sector with the conserved current it vanishes, thereby it has no dynamics and therefore we can throw it away. On the other hand the transversal sector can be re-written in such a way that,

$$\langle B^\mu B^\nu \rangle_T(k) = \frac{i}{(k^2 - \rho_+^2)(k^2 - \rho_-^2)} \left[(k^2 - \alpha^2) \left(\eta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right) + 2mi\epsilon^{\mu\lambda\nu} k_\lambda \right], \quad (37)$$

where the massive poles are,

$$\rho_\pm^2 = \alpha^2 + 2m^2 \pm 2m\sqrt{\alpha^2 + m^2}. \quad (38)$$

² It is different from what occurs to the solutions of the scalar case of Nielsen-Olesen vortex, which are dependent on a real parameter that comes from a breaking of the internal symmetry $U(1)$.

We can see that the transversal sector poles represent the unique dynamic degrees of freedom of the model, and if we take $m = 0$ we get the usual Proca-type massive pole as expected. The important point here is that we can easily observe that the topological mass contributes to dynamic mass poles $\rho_{\pm}^2$. And to obtain such contribution we suggest a simple relation between the mass and the topological mass terms, that is

$$\varepsilon = \frac{m}{\alpha}, \quad (39)$$

which means that the topological mass is a multiple of the usual mass term, and ε is a constant value. In this way we can substitute it in (38), giving

$$\rho_{\pm}^2 = C(\varepsilon)\alpha^2, \quad (40)$$

where $C(\varepsilon)$ is the constant expression

$$C(\varepsilon) = 1 + 2\varepsilon^2 \pm 2\varepsilon(1 + \varepsilon^2)^{1/2}. \quad (41)$$

We observe that for real values $C(\varepsilon)$ is always positive.

Therefore we can conclude that the contribution of the topological sector does not spoil the SSB will be procedure in the sense that if we take the mass-term α^2 to be negative thus will be the Chern-Simons-type mass term m^2 . However, in this approach, the Chern-Simons-type sector does not play a role in the vacuum computation, it indicates the “speed” of the field in order to arrive at the asymptotic limit and the size of the region where H is different from zero or the vortex core. So, in order to include a Chern-Simons-type contribution to the mass pole we have to “deform” the mass term in the above $C(\varepsilon)$ in such way that

$$\alpha^2 \rightarrow C(\varepsilon)\alpha^2 \quad (42)$$

where $C(\varepsilon) \geq 1$ and it is easy to see that for $m = 0$ implies that $\varepsilon = 0$ and $C(\varepsilon) = 1$. So the vector solitons can have two modes ($\pm$) that depend on the mass deformation factor $C(\varepsilon)$, noticing that they are both positive.

Therefore, taking the asymptotic behavior of the field and neglecting the $\pm$ indexes, we can verify that the function (32) is consistent with the physical limits defined in (21) and (22). We also notice that this definition can be extended to the following quantities,

$$\beta_{\mu}(r)\beta^{\mu}(r) = |\beta(r)|^2 = b_{\mu}b^{\mu} \left(1 - ae^{-\sqrt{-\rho^2}r}\right)^2, \quad (43)$$

which we can take,

$$|\beta(r)| = |b| \left(1 - ae^{-\sqrt{-\rho^2}r}\right), \quad (44)$$

and taking the $r \rightarrow \infty$ limit we have that $|\beta(r)|_{r \rightarrow \infty} = |b|$ which is consistent with the definitions adopted. This result implies that each component of $\beta_{\mu}(r)$ behaves as a unique soliton traveling in various directions, and whose distance is related to the mass parameter ρ^2 . The vector character of the soliton is the consequence of the solitonic modes³ in the $(1+2)D$ space-time.

The non-trivial vacuum value expressed in the relation (32) suggests that the vector soliton is generated from the Lorentz symmetry violation represented by the b_{μ} parameter. Furthermore, vector solitons are also associated with local anisotropy of the vacuum energy (or optical vacuum) which can be obtained from the non-trivial vacuum state of vector matter field model, as has been speculated in a previous work[26]. For this reason, we can notice that vector soliton objects which present independent components of polarization b_{μ} and originate from Lorentz symmetry violation are indeed not a simple extension of scalar topological solitons.

In order to obtain the factors of the function (44) a further analysis of $\beta_{\mu}(r)$ beyond of these limits can be made. To this end we suggest that the factor a can be a function of r in such a way that it has to satisfy $\lim_{r \rightarrow 0} \beta_{\mu}(r) = 0$. The best form of the factor a is given by,

$$a(r) = \frac{\sin\left(\sqrt{-\rho^2}r\right)}{\sqrt{-\rho^2}r}, \quad (45)$$

³ It can be related to collective behavior of these solitonic modes.




FIG. 1: Plot of $\beta(r)$ field versus the distance r given by expression (46). The mass parameter ρ is estimated as the Higgs mass value. We observe that $\beta(r)$ assumes asymptotic values for $|b| = 1$ at infinity.

so, substituting this expression in the function (44), we have that

$$\beta_\mu(r) = b_\mu \left(1 - \frac{\sin(\sqrt{-\rho^2} r)}{\sqrt{-\rho^2} r} e^{-\sqrt{-\rho^2} r} \right), \quad (46)$$

where we verify that $\beta_\mu(r)$ is a smooth and analytic function for all real values of r , and it satisfies both limits expressed in (21). Then the expression (46) might be taken as a simplified analytical solution for the function $\beta(r)$ near to the asymptotic limit. We can observe in Fig. 1, that it fits very well, particularly for values of ρ to the order of Higgs mass. From Fig. 1 we can also notice a smooth convergence of $\beta(r)$ at the asymptotic limit or $|b| = 1$. The vector $B_\mu(r)$ shows a spontaneous symmetry breaking due to the parameters b_μ present in each component. This suggests that a “multi-soliton” with a vector field structure could be generated.

V. CONCLUSION

In this work we have analyzed the contribution of the non-trivial vacuum to the topological behavior of a vector matter field in $(1 + 2)D$. In spite of a non-exact and/or non-analytic solution of the differential equations derived from each component of the equation of motion the topological characteristic of the solution is guaranteed. In the present model, we have obtained that the charged vortices show similar behavior as in the Maxwell-Chern-Simons model[27, 28]. Moreover, the resulting topological objects could be interpreted as vector solitons, where each one of their components are shown to be a polarized soliton in a spacetime direction. We have also shown that a possible origin for these vector solitons comes from the non-trivial vacuum in the vector matter field B_μ dynamics. Furthermore we have analyzed the role played by the topological Chern-Simons-type terms. Although they do not contribute to the non-trivial vacuum obtained, by means of a mass pole verification using propagators computation we observe that it can contribute with a positive two-mode-extension to the mass term, which indicates that its factor preserves the topological feature of the asymptotic limit of the field. It can characterize the size of the region where the “magnetic” field is non-zero, or core vortex, and the “speed” of the field to saturate the asymptotic limit.

In the present work we have noticed that the soliton modes with polarized direction b_μ generated can be associated with local anisotropy of vacuum energy. There is an interesting similarity to the study of the Lorentz symmetry violation studies [14, 15, 16, 17, 18, 19]. In this work, the (local) Lorentz symmetry violation might induce a local spacetime perturbation giving a dynamical anisotropic anomaly that is manifest by a vector soliton object. Indeed, it suggests the possible existence of topological vector solitons in the limit of very high energy physics.

We can also speculate that the vector solitons described in this work could be connected to the multi-soliton modes in the context of nonlinear crystals, which first appear in the work of Manakov for low energy systems[29]. We also noticed that such multi-solitons (or solitons with many components) were recently observed in optical fiber experiments[30, 31, 32]. From another point of view, experiments have been performed to verify the presence of

non-topological multi-solitons in anisotropic crystals in $(1+2)D$ [33, 34] that could be correlated with the present work.

In the scope of the Quantum Field Theory it could be noted that topological vector solitons stem from a possible optical vacuum energy state or dynamical topological defect. Thus these objects might be present in models that due to the space structure have anisotropic (optical or not) effects.

VI. ACKNOWLEDGMENTS

LPC would like to thank M.T.D. Orlando for his kind hospitality at the Physics Department of the Universidade Federal do Espírito Santo (UFES) and to Prof. J.A. Helay el-Neto for the kind hospitality at the CBPF. The authors would like to thank Susan Harris de Mello for the careful reading and the suggestions to improve the manuscript.

-
- [1] H.S. Snyder, *Phys. Rev.* **71**, 38(1947).
 - [2] C.N. Yang, *Phys. Rev.* **72**, 874(1947).
 - [3] M.R. Douglas, N.A. Nekrasov, *Rev. Mod. Phys.* **73**, 977(2001).
 - [4] N. Seiberg, E. Witten, *J.H.E.P.* **9909**, 032(1999).
 - [5] R. Jackiw, *Nucl. Phys. Proc. Suppl.* **108**, 30(2002).
 - [6] D. Bahns, S. Doplicher, K. Fredenhagen, G. Piacitelli, *Phys. Lett.* **B533**, 178(2000).
 - [7] J.L. Hewett, F.J. Petriello, T.G. Rizzo, *Phys. Rev.* **D64**, 075012(2001).
 - [8] I. Mocioiu, M. Pospelov, R. Roiban, *Phys. Lett.* **B489**, 390(2000).
 - [9] C.E. Carlson, C.D. Carone, R.F. Lebed, *Phys. Lett.* **B518**, 201(2001).
 - [10] Xavier Calmet, *Phys. Rev.* **D71**, 085012 (2005).
 - [11] G. Berrino, S.L. Cacciatori, A. Celi, L. Martucci, A. Vicini, *Phys. Rev.* **D67**, 065021 (2002).
 - [12] V.O. Rivelles, *Proc. Sci.* **WC2004**, 029(2004).
 - [13] G. Moulataka, M. Rausch de Trautenberg, A. Tanasa, *Non-trivial extension of the Poincare algebra for antisymmetric gauge fields*, 11th International Conference on Symmetry Methods in Physics (SYMPHYS-11), Prague, Czech Republic, 21-24 Jun 2004, hep-th/0407168;
N. Mohammadi, G. Moulataka, M. Rausch de Trautenberg, *Int. J. Mod. Phys.* **A19** (2004) 5585;
M. Rausch de Trautenberg, M.J. Slupinski, *J. Math. Phys.* **41** (2000) 4556;
M. Rausch de Trautenberg, M.J. Slupinski, *Mod. Phys. Lett.* **A12** (1997) 3051;
L.P. Colatto, J.L. Matheus-Valle, *J. Math. Phys.* **37** (1996) 6121.
 - [14] D. Colladay, V.A. Kosteleck y, *Phys. Rev.* **D55**, 6760 (1997).
 - [15] D. Colladay, V.A. Kosteleck y, *Phys. Rev.* **D58**, 116002 (1998).
 - [16] G. Amelino-Camelia, *The three perspectives on the quantum-gravity problem and their implications for the fate of Lorentz symmetry*, gr-qc/0309054;
O. Bertolami, R. Lehnert, A. Ribeiro, *Phys. Rev.* **D69**, 083513 (2004);
O. Bertolami, *Lect. Notes Phys.* **633**, 96 (2003);
R. Lehnert, *Phys. Rev.* **D68**, 085003 (2003);
V. A. Kostelecky, R. Lehnert, M. J. Perry, *Phys. Rev.* **D68**, 123511 (2003);
E.C. Vagenas, *J.H.E.P.* **0307**, 046 (2003);
J. W. Moffat, *Int. J. Mod. Phys.* **D12**, 1279 (2003);
S. Das, G. Kunstatter, *Class. Quant. Grav.* **20**, 2015 (2003);
J.D. Barrow, *Phys. Lett.* **B564**, 1 (2003);
P.C.W. Davies, T.M. Davis, C.H. Lineweaver, *Nature* **418**, 602 (2002);
V. A. Kostelecky, R. Lehnert, *Phys. Rev.* **D63**, 065008 (2001);
K. Greisen, *Phys. Rev. Lett.* **16**, 748 (1966);
G.T. Zatsepin, V.A. Kuzmin, *Pisma Zh. Eksp. Teor. Fiz.* **4**, 114 (1966).
 - [17] A.A. Andrianov, R. Soldati, *Phys. Rev.* **D51**, 5961 (1995).
 - [18] A.A. Andrianov, R. Soldati, *Phys. Lett.* **B435**, 449 (1998).
 - [19] I.V. Bar'ykhtar, B.A. Ivanov, *Phys. Lett.* **A98**, 222(1983).
 - [20] S. Carrol, G. Field, R. Jackiw, *Phys. Rev.* **D41**, 1231 (1999).
 - [21] D. Hutsem ekers, H. Lamy, *Astron. Astrophys.* **367**, 381 (2001);
D. Hutsem ekers and H. Lamy, *Astron. Astrophys.* **332**, 410 (1998);
M. Goldhaber, V. Trimble, *J. Astrophys. Astron.* **17**, 17 (1996).
 - [22] H.B. Nielsen, I. Picek *Nucl. Phys.* **B211**, 269(1983).
 - [23] D. Harari, P. Sikivie, *Phys. Lett.* **B289**, 67(1992).
 - [24] B. Nodland, V.A. Ralston, *Phys. Rev. Lett.* **78**, 3043(1997).
 - [25] B. Nodland, V.A. Ralston, *Phys. Rev. Lett.* **79**, 1958(1997).

- [26] L.P. Colatto, A.L.A. Penna, W.C. Santos, *Eur. Phys. J.* **C36**, 79(2004).
- [27] L.H. Ryder, *Quantum Field Theory*, second edition, Cambridge University Press (1996).
- [28] L.P. Colatto, A.L.A. Penna, W.C. Santos, *in preparation*.
- [29] S.V. Manakov, *J.E.T.P.* **38**, 3043(1974).
- [30] J.U. Kang, G.I. Stegeman, J.S. Aichinson, N. Akhmediev, *Phys. Rev. Lett.* **76**, 3699(1996).
- [31] Y. Barat, Y. Silberberg, *Phys. Rev. Lett.* **78**, 3290(1997).
- [32] M. Seveg, M. Shin, G.C. Valley *J. Opt. Soc. Am.* **B13**, 706(1995).
- [33] D.N. Christodoulides, M.I. Carvalho, *J. Opt. Soc. Am.* **B12**, 1628(1995).
- [34] D.N. Christodoulides, M.I. Carvalho, *Opt. Lett.* **19**, 1714(1994).